

SMC3 Rabbit mAb

Catalog # AP77236

Product Information

Application	WB, IHC-P, IF, FC, ICC
Primary Accession	Q9UQE7
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human SMC3
Purification	Affinity Chromatography
Calculated MW	141542

Additional Information


Gene ID	9126
Other Names	SMC3
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	SMC3
Synonyms	BAM, BMH, CSPG6, SMC3L1
Function	Central component of cohesin, a complex required for chromosome cohesion during the cell cycle. The cohesin complex may form a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. Cohesion is coupled to DNA replication and is involved in DNA repair. The cohesin complex also plays an important role in spindle pole assembly during mitosis and in chromosomes movement.
Cellular Location	Nucleus {ECO:0000250 UniProtKB:Q9CW03}. Chromosome {ECO:0000250 UniProtKB:Q9CW03}. Chromosome, centromere {ECO:0000250 UniProtKB:Q9CW03}. Note=Associates with chromatin. Before prophase it is scattered along chromosome arms. During prophase, most of

cohesin complexes dissociate from chromatin probably because of phosphorylation by PLK, except at centromeres, where cohesin complexes remain. At anaphase, the RAD21 subunit of the cohesin complex is cleaved, leading to the dissociation of the complex from chromosomes, allowing chromosome separation. The phosphorylated form at Ser-1083 is preferentially associated with unsynapsed chromosomal regions (By similarity). {ECO:0000250|UniProtKB:Q9CW03}

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.