

# TXK Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7723a

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P42681</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB03245
<b>Calculated MW</b>	61258
<b>Antigen Region</b>	22-52

## Additional Information

---

<b>Gene ID</b>	7294
<b>Other Names</b>	Tyrosine-protein kinase TXK, Protein-tyrosine kinase 4, Resting lymphocyte kinase, TXK, PTK4, RLK
<b>Target/Specificity</b>	This TXK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 22-52 amino acids from the N-terminal region of human TXK.
<b>Dilution</b>	WB~~1:2000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TXK Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	TXK
<b>Synonyms</b>	PTK4, RLK
<b>Function</b>	Non-receptor tyrosine kinase that plays a redundant role with ITK in regulation of the adaptive immune response. Regulates the development,

function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation leads to the recruitment of TXK to the cell membrane, where it is phosphorylated at Tyr-420. Phosphorylation leads to TXK full activation. Also contributes to signaling from many receptors and participates in multiple downstream pathways, including regulation of the actin cytoskeleton. Like ITK, can phosphorylate PLCG1, leading to its localization in lipid rafts and activation, followed by subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. Plays a role in the positive regulation of IFNG transcription in T- helper 1 cells as part of an IFNG promoter-binding complex with PARP1 and EEF1A1 (PubMed:[11859127](#), PubMed:[17177976](#)). Within the complex, phosphorylates both PARP1 and EEF1A1 (PubMed:[17177976](#)). Also phosphorylates key sites in LCP2 leading to the up-regulation of Th1 preferred cytokine IL-2. Phosphorylates 'Tyr-201' of CTLA4 which leads to the association of PI-3 kinase with the CTLA4 receptor.

#### Cellular Location

Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein. Note=Localizes in the vicinity of cell surface receptors in the plasma membrane after receptor stimulation Translocates into the nucleus and enhances IFN-gamma gene transcription in T-cells

#### Tissue Location

Expressed in T-cells and some myeloid cell lines. Expressed in Th1/Th0 cells with IFN-gamma-producing potential

## Background

---

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the  $\gamma$  phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway.

## References


---

Takeba, Y., et al., J. Immunol. 168(5):2365-2370 (2002). Spritz, R.A., et al., Genomics 22(2):431-436 (1994). Haire, R.N., et al., Hum. Mol. Genet. 3(6):897-901 (1994).



## Images

---

Western blot analysis of TXK (arrow) using rabbit polyclonal hTXK-E37 (Cat. #AP7723a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the TXK gene.



Western blot analysis of anti-TXK Pab (Cat. #AP7723a) in Jurkat cell lysate. TXK (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.