

PAX5 Rabbit mAb

Catalog # AP77263

Product Information

Application WB, IHC-P, IF, FC, ICC

Primary Accession <u>Q02548</u>

Reactivity Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human PAX5

Purification Affinity Chromatography

Calculated MW 42149

Additional Information

Gene ID 5079

Other Names PAX5

Dilution WB~~1/500-1/1000 IHC-P~~N/A IF~~1/50-1/200 FC~~1:10~50 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name PAX5

Function Transcription factor that plays an essential role in commitment of lymphoid

progenitors to the B-lymphocyte lineage (PubMed: 10811620,


PubMed: 27181361). Fulfills a dual role by repressing B-lineage inappropriate


genes and simultaneously activating B-lineage- specific genes


(PubMed:<u>10811620</u>, PubMed:<u>27181361</u>). In turn, regulates cell adhesion and migration, induces V(H)-to-D(H)J(H) recombination, facilitates pre-B-cell receptor signaling and promotes development to the mature B-cell stage (PubMed:<u>32612238</u>). Repression of the cohesin- release factor WAPL causes global changes of the chromosomal architecture in pro-B cells to facilitate the

generation of a diverse antibody repertoire (PubMed:32612238).

Cellular Location Nucleus.







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.