

ERK5 Rabbit mAb

Catalog # AP77285

Product Information

Application	WB, FC, IP, ICC
Primary Accession	Q13164
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	88386


Additional Information

Gene ID	5598
Other Names	MAPK7
Dilution	WB~~1/500-1/1000 FC~~1:10~50 IP~~N/A ICC~~N/A
Format	Liquid

Protein Information

Name	MAPK7
Synonyms	BMK1, ERK5, PRKM7
Function	<p>Plays a role in various cellular processes such as proliferation, differentiation and cell survival. The upstream activator of MAPK7 is the MAPK kinase MAP2K5. Upon activation, it translocates to the nucleus and phosphorylates various downstream targets including MEF2C. EGF activates MAPK7 through a Ras-independent and MAP2K5-dependent pathway. As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via interaction with STUB1/CHIP and promotion of STUB1-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity). May have a role in muscle cell differentiation. May be important for endothelial function and maintenance of blood vessel integrity. MAP2K5 and MAPK7 interact specifically with one another and not with MEK1/ERK1 or MEK2/ERK2 pathways. Phosphorylates SGK1 at Ser-78 and this is required for growth factor-induced cell cycle progression. Involved in the regulation of p53/TP53 by disrupting the PML-MDM2 interaction.</p>
Cellular Location	Cytoplasm. Nucleus. Nucleus, PML body. Note=Translocates to the nucleus upon activation
Tissue Location	Expressed in many adult tissues. Abundant in heart, placenta, lung, kidney and skeletal muscle. Not detectable in liver

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.