

Caspase 5 Rabbit mAb

Catalog # AP77305

Product Information

Application WB, IHC-P, IF, ICC, IP

Primary Accession P51878
Reactivity Human
Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human Caspase 5

Purification Affinity Chromatography

Calculated MW 49736

Additional Information

Gene ID 838

Other Names CASP5

Dilution WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A IP~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name CASP5 {ECO:0000303 | PubMed:16893518, ECO:0000312 | HGNC:HGNC:1506}

Function Thiol protease that acts as a mediator of programmed cell death

(PubMed:<u>28314590</u>, PubMed:<u>29898893</u>). Initiates pyroptosis, a programmed lytic cell death pathway through cleavage of Gasdermin-D (GSDMD): cleavage releases the N-terminal gasdermin moiety (Gasdermin-D, N-terminal) that

binds to membranes and forms pores, triggering pyroptosis


(PubMed:<u>29898893</u>). Also mediates cleavage and maturation of IL18 (PubMed:<u>37993714</u>). Cleavage of GSDMD and IL18 is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP4 (PubMed:<u>37993714</u>). During non-canonical inflammasome activation, cuts CGAS and may play a role in the regulation of antiviral innate immune


activation (PubMed:28314590).

Tissue Location Expressed in barely detectable amounts in most tissues except brain, highest

levels being found in lung, liver and skeletal muscle.

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.