

RPS3 Rabbit mAb

Catalog # AP77331

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	P23396
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human RPS3
Purification	Affinity Chromatography
Calculated MW	26688

Additional Information

Gene ID	6188
Other Names	RPS3
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information



Name	RPS3 {ECO:0000303 PubMed:11875025}
Function	Component of the small ribosomal subunit (PubMed: 23636399 , PubMed: 8706699). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed: 23636399 , PubMed: 8706699). Has endonuclease activity and plays a role in repair of damaged DNA (PubMed: 7775413). Cleaves phosphodiester bonds of DNAs containing altered bases with broad specificity and cleaves supercoiled DNA more efficiently than relaxed DNA (PubMed: 15707971). Displays high binding affinity for 7,8-dihydro-8-oxoguanine (8-oxoG), a common DNA lesion caused by reactive oxygen species (ROS) (PubMed: 14706345). Has also been shown to bind with similar affinity to intact and damaged DNA (PubMed: 18610840). Stimulates the N-glycosylase activity of the base excision protein OGG1 (PubMed: 15518571). Enhances the uracil excision activity of UNG1 (PubMed: 18973764). Also stimulates the cleavage of the phosphodiester backbone by APEX1 (PubMed: 18973764). When located in the mitochondrion,

reduces cellular ROS levels and mitochondrial DNA damage (PubMed:[23911537](#)). Has also been shown to negatively regulate DNA repair in cells exposed to hydrogen peroxide (PubMed:[17049931](#)). Plays a role in regulating transcription as part of the NF-kappa-B p65-p50 complex where it binds to the RELA/p65 subunit, enhances binding of the complex to DNA and promotes transcription of target genes (PubMed:[18045535](#)). Represses its own translation by binding to its cognate mRNA (PubMed:[20217897](#)). Binds to and protects TP53/p53 from MDM2-mediated ubiquitination (PubMed:[19656744](#)). Involved in spindle formation and chromosome movement during mitosis by regulating microtubule polymerization (PubMed:[23131551](#)). Involved in induction of apoptosis through its role in activation of CASP8 (PubMed:[14988002](#)). Induces neuronal apoptosis by interacting with the E2F1 transcription factor and acting synergistically with it to up-regulate pro-apoptotic proteins BCL2L1/BIM and HRK/Dp5 (PubMed:[20605787](#)). Interacts with TRADD following exposure to UV radiation and induces apoptosis by caspase-dependent JNK activation (PubMed:[22510408](#)).

Cellular Location

Cytoplasm. Nucleus. Nucleus, nucleolus Mitochondrion inner membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, spindle. Note=In normal cells, located mainly in the cytoplasm with small amounts in the nucleus but translocates to the nucleus in cells undergoing apoptosis (By similarity). Nuclear translocation is induced by DNA damaging agents such as hydrogen peroxide (PubMed:[17560175](#)). Accumulates in the mitochondrion in response to increased ROS levels (PubMed:[23911537](#)) Localizes to the spindle during mitosis (PubMed:[23131551](#)). Localized in cytoplasmic mRNP granules containing untranslated mRNAs (PubMed:[17289661](#)).
{ECO:0000250|UniProtKB:P62908, ECO:0000269|PubMed:[17289661](#), ECO:0000269|PubMed:[17560175](#), ECO:0000269|PubMed:[23131551](#), ECO:0000269|PubMed:[23911537](#)}

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.