

NGFR Rabbit mAb

Catalog # AP77343

Product Information

Application	WB, IHC-P, IP, ICC
Primary Accession	P08138
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	45183



Additional Information

Gene ID	4804
Other Names	NGFR
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A ICC~~N/A
Format	Liquid

Protein Information

Name	NGFR
Synonyms	TNFRSF16
Function	<p>Low affinity receptor which can bind to NGF, BDNF, NTF3, and NTF4. Forms a heterodimeric receptor with SORCS2 that binds the precursor forms of NGF, BDNF and NTF3 with high affinity, and has much lower affinity for mature NGF and BDNF (PubMed:24908487). Plays an important role in differentiation and survival of specific neuronal populations during development (By similarity). Can mediate cell survival as well as cell death of neural cells. Plays a role in the inactivation of RHOA (PubMed:26646181). Plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin- dependent glucose uptake (By similarity). Necessary for the circadian oscillation of the clock genes BMAL1, PER1, PER2 and NR1D1 in the suprachiasmatic nucleus (SCMgetaN) of the brain and in liver and of the genes involved in glucose and lipid metabolism in the liver (PubMed:23785138). Together with BFAR negatively regulates NF-kappa-B and JNK-related signaling pathways (PubMed:22566094).</p>
Cellular Location	<p>Cell membrane; Single-pass type I membrane protein. Cytoplasm. Perikaryon {ECO:0000250 UniProtKB:Q9Z0W1}. Cell projection, growth cone {ECO:0000250 UniProtKB:Q9Z0W1}. Cell projection, dendritic spine</p>

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.