

PUM1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7734a

Product Information

Application	WB, IHC-P, E
Primary Accession	Q14671
Other Accession	Q80U78
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB10535
Calculated MW	126473
Antigen Region	1-30

Additional Information

Gene ID	9698
Other Names	Pumilio homolog 1, HsPUM, Pumilio-1, PUM1, KIAA0099, PUMH1
Target/Specificity	This PUM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human PUM1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PUM1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PUM1 (HGNC:14957)
Function	Sequence-specific RNA-binding protein that acts as a post- transcriptional repressor by binding the 3'-UTR of mRNA targets. Binds to an RNA consensus sequence, the Pumilio Response Element (PRE), 5'- UGUANAUA-3', that is

related to the Nanos Response Element (NRE) (PubMed:[18328718](#), PubMed:[21397187](#), PubMed:[21572425](#), PubMed:[21653694](#)). Mediates post-transcriptional repression of transcripts via different mechanisms: acts via direct recruitment of the CCR4-POP2-NOT deadenylase leading to translational inhibition and mRNA degradation (PubMed:[22955276](#)). Also mediates deadenylation-independent repression by promoting accessibility of miRNAs (PubMed:[18776931](#), PubMed:[20818387](#), PubMed:[20860814](#), PubMed:[22345517](#)). Following growth factor stimulation, phosphorylated and binds to the 3'-UTR of CDKN1B/p27 mRNA, inducing a local conformational change that exposes miRNA-binding sites, promoting association of miR-221 and miR-222, efficient suppression of CDKN1B/p27 expression, and rapid entry to the cell cycle (PubMed:[20818387](#)). Acts as a post-transcriptional repressor of E2F3 mRNAs by binding to its 3'- UTR and facilitating miRNA regulation (PubMed:[22345517](#), PubMed:[29474920](#)). Represses a program of genes necessary to maintain genomic stability such as key mitotic, DNA repair and DNA replication factors. Its ability to repress those target mRNAs is regulated by the lncRNA NORAD (non-coding RNA activated by DNA damage) which, due to its high abundance and multitude of PUMILIO binding sites, is able to sequester a significant fraction of PUM1 and PUM2 in the cytoplasm (PubMed:[26724866](#)). Involved in neuronal functions by regulating ATXN1 mRNA levels: acts by binding to the 3'-UTR of ATXN1 transcripts, leading to their down-regulation independently of the miRNA machinery (PubMed:[25768905](#), PubMed:[29474920](#)). Plays a role in cytoplasmic sensing of viral infection (PubMed:[25340845](#)). In testis, acts as a post- transcriptional regulator of spermatogenesis by binding to the 3'-UTR of mRNAs coding for regulators of p53/TP53. Involved in embryonic stem cell renewal by facilitating the exit from the ground state: acts by targeting mRNAs coding for naive pluripotency transcription factors and accelerates their down-regulation at the onset of differentiation (By similarity). Binds specifically to miRNA MIR199A precursor, with PUM2, regulates miRNA MIR199A expression at a postranscriptional level (PubMed:[28431233](#)).

Cellular Location

Cytoplasm. Cytoplasm, P-body. Cytoplasmic granule. Note=Recruited to cytoplasmic stress granules upon viral infection.

Tissue Location

Expressed in brain, heart, kidney, muscle, intestine and stomach. Not expressed in cerebellum, corpus callosum, caudate nucleus, hippocampus, medulla oblongata and putamen. Expressed in all fetal tissues tested.

Background



PUM1 is a sequence-specific RNA-binding protein that regulates translation and mRNA stability by binding the 3'-UTR of mRNA targets. Its interactions and tissue specificity suggest that it may be required to support proliferation and self-renewal of stem cells by regulating the translation of key transcripts.

References

- Kusz,K., Mol. Reprod. Dev. 74 (6), 795-799 (2007)
 Spik,A., Reprod Biol 6 SUPPL 1, 37-42 (2006)
 Moore,F.L., Proc. Natl. Acad. Sci. U.S.A. 100 (2), 538-543 (2003)

Images

Western blot analysis of PUM1 (Human N-term) (Cat. #AP7734a) in HL-60 cell line lysates (35ug/lane). PUM1 (arrow) was detected using the purified Pab.(2ug/ml)



Formalin-fixed and paraffin-embedded human testis tissue reacted with PUM1 Antibody (N-term) (Cat.#AP7734a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.