

NRAS Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7745b

Product Information

Application	WB, E
Primary Accession	P01111
Other Accession	Q2MJK3
Reactivity	Human, Rat, Mouse
Predicted	Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21229
Antigen Region	147-179

Additional Information

Gene ID	4893
Other Names	GTPase NRas, Transforming protein N-Ras, NRAS, HRAS1
Target/Specificity	This NRAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 147-179 amino acids from the C-terminal region of human NRAS.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NRAS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NRAS
Synonyms	HRAS1
Function	Ras proteins bind GDP/GTP and possess intrinsic GTPase activity.

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor Note=Shuttles between the plasma membrane and the Golgi apparatus

Background


NRAS is a membrane protein that shuttles between the Golgi apparatus and the plasma membrane. This shuttling is regulated through palmitoylation and depalmitoylation by the ZDHHC9-GOLGA7 complex. This protein, which has intrinsic GTPase activity, is activated to a GTP-bound form by a GTPase activating protein and inactivated to a GDP-bound form by a guanine nucleotide-exchange factor. Defects in the gene encoding this protein are a cause of juvenile myelomonocytic leukemia (JMML).

References

Smalley,K.S., Cancer Res. 68 (14), 5743-5752 (2008)

Banerji,U., Mol. Cancer Ther. 7 (4), 737-739 (2008)

Images



Anti-NRAS Antibody (C-term) at 1:2000 dilution + MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.