

Dysferlin Rabbit mAb

Catalog # AP77483

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	O75923
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Dysferlin
Purification	Affinity Chromatography
Calculated MW	237295

Additional Information

Gene ID	8291
Other Names	DYSF
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	DYSF
Synonyms	FER1L1
Function	Key calcium ion sensor involved in the Ca(2+)-triggered synaptic vesicle-plasma membrane fusion. Plays a role in the sarcolemma repair mechanism of both skeletal muscle and cardiomyocytes that permits rapid resealing of membranes disrupted by mechanical stress (By similarity).
Cellular Location	Cell membrane, sarcolemma; Single-pass type IV membrane protein. Cytoplasmic vesicle membrane {ECO:0000250 UniProtKB:Q9ESD7}; Single-pass type IV membrane protein. Cell membrane; Single-pass type IV membrane protein. Late endosome membrane; Single-pass type IV membrane protein. Note=Colocalizes, during muscle differentiation, with BIN1 in the T-tubule system of myotubules and at the site of contact between two myotubes or a myoblast and a myotube. Wounding of myotubes led to its

focal enrichment to the site of injury and to its relocalization in a Ca²⁺-dependent manner toward the plasma membrane. Colocalizes with AHNAK, AHNAK2 and PARVB at the sarcolemma of skeletal muscle. Detected on the apical plasma membrane of the syncytiotrophoblast. Reaches the plasma membrane through a caveolin-independent mechanism. Retained by caveolin at the plasma membrane (By similarity). Colocalizes, during muscle differentiation, with CACNA1S in the T-tubule system of myotubules (By similarity). Accumulates and colocalizes with fusion vesicles at the sarcolemma disruption sites (By similarity) {ECO:0000250|UniProtKB:Q9ESD7}

Tissue Location

Expressed in skeletal muscle, myoblast, myotube and in the syncytiotrophoblast (STB) of the placenta (at protein level) Ubiquitous. Highly expressed in skeletal muscle. Also found in heart, brain, spleen, intestine, placenta and at lower levels in liver, lung, kidney and pancreas.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.