

# TIEG Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7749a

## Product Information

---

Application	WB, IHC-P, E
Primary Accession	<a href="#">Q13118</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB14876
Calculated MW	52555
Antigen Region	13-40

## Additional Information

---

Gene ID	7071
Other Names	Krueppel-like factor 10, EGR-alpha, Transforming growth factor-beta-inducible early growth response protein 1, TGFB-inducible early growth response protein 1, TIEG-1, KLF10, TIEG, TIEG1
Target/Specificity	This TIEG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 13-40 amino acids from the N-terminal region of human TIEG.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TIEG Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	KLF10
Synonyms	TIEG, TIEG1
Function	Transcriptional repressor which binds to the consensus sequence

5'-GGTGTG-3'. Plays a role in the regulation of the circadian clock; binds to the GC box sequence in the promoter of the core clock component ARTNL/BMAL1 and represses its transcriptional activity. Regulates the circadian expression of genes involved in lipogenesis, gluconeogenesis, and glycolysis in the liver. Represses the expression of PCK2, a rate-limiting step enzyme of gluconeogenesis (By similarity). May play a role in the cell cycle regulation.

## Cellular Location

Nucleus {ECO:0000250|UniProtKB:O89091}.


## Background

TIEG is a transcriptional repressor involved in the regulation of cell growth. TIEG binds to the consensus sequence 5'-GGTGTG-3' and inhibits cell growth.


## References

Hawse,J.R., Mol. Endocrinol. 22 (7), 1579-1595 (2008)  
 Ivanov,S.V., Biochem. Biophys. Res. Commun. 370 (4), 536-540 (2008)  
 Jin,W., FEBS Lett. 581 (20), 3826-3832 (2007)  
 Fueki,N., Respiration 74 (4), 454-459 (2007)

## Images



All lanes : Anti-TIEG Antibody (N-term) at 1:2000 dilution  
 Lane 1: 293T whole cell lysates Lane 2: MCF-7 whole cell lysates Lane 3: human kidney lysates Lane 4: human lung lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 53 kDa  
 Blocking/Dilution buffer: 5% NFDm/TBST.



Formalin-fixed and paraffin-embedded human prostate carcinoma tissue reacted with TIEG antibody (N-term) (Cat.#AP7749a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.