

FOXP3 Rabbit mAb

Catalog # AP77586

Product Information

Application	WB, IHC-P
Primary Accession	Q9BZS1
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human FOXP3
Purification	Affinity Chromatography
Calculated MW	47244

Additional Information

Gene ID	50943
Other Names	FOXP3
Dilution	WB~~1/500-1/1000 IHC-P~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	FOXP3
Synonyms	IPEX
Function	Transcriptional regulator which is crucial for the development and inhibitory function of regulatory T-cells (Treg) (PubMed: 17377532 , PubMed: 21458306 , PubMed: 23947341 , PubMed: 24354325 , PubMed: 24722479 , PubMed: 24835996 , PubMed: 30513302 , PubMed: 32644293). Plays an essential role in maintaining homeostasis of the immune system by allowing the acquisition of full suppressive function and stability of the Treg lineage, and by directly modulating the expansion and function of conventional T-cells (PubMed: 23169781). Can act either as a transcriptional repressor or a transcriptional activator depending on its interactions with other transcription factors, histone acetylases and deacetylases (PubMed: 17377532 , PubMed: 21458306 , PubMed: 23947341 , PubMed: 24354325 , PubMed: 24722479). The suppressive activity of Treg involves the coordinate

activation of many genes, including CTLA4 and TNFRSF18 by FOXP3 along with repression of genes encoding cytokines such as interleukin-2 (IL2) and interferon-gamma (IFNG) (PubMed:[17377532](#), PubMed:[21458306](#), PubMed:[23947341](#), PubMed:[24354325](#), PubMed:[24722479](#)). Inhibits cytokine production and T-cell effector function by repressing the activity of two key transcription factors, RELA and NFATC2 (PubMed:[15790681](#)). Mediates transcriptional repression of IL2 via its association with histone acetylase KAT5 and histone deacetylase HDAC7 (PubMed:[17360565](#)). Can activate the expression of TNFRSF18, IL2RA and CTLA4 and repress the expression of IL2 and IFNG via its association with transcription factor RUNX1 (PubMed:[17377532](#)). Inhibits the differentiation of IL17 producing helper T-cells (Th17) by antagonizing RORC function, leading to down-regulation of IL17 expression, favoring Treg development (PubMed:[18368049](#)). Inhibits the transcriptional activator activity of RORA (PubMed:[18354202](#)). Can repress the expression of IL2 and IFNG via its association with transcription factor IKZF4 (By similarity).

Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00089, ECO:0000269 | PubMed:17360565, ECO:0000269 | PubMed:18354202, ECO:0000269 | PubMed:22678915, ECO:0000269 | PubMed:23396208, ECO:0000269 | PubMed:23973222, ECO:0000269 | PubMed:23973223, ECO:0000269 | PubMed:32644293}. Cytoplasm Note=Predominantly expressed in the cytoplasm in activated conventional T-cells whereas predominantly expressed in the nucleus in regulatory T- cells (Treg). The 41 kDa form derived by proteolytic processing is found exclusively in the chromatin fraction of activated Treg cells (By similarity). {ECO:0000250 | UniProtKB:Q99JB6, ECO:0000269 | PubMed:22678915}

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.