

# GHRH Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7758A

## Product Information

---

<b>Application</b>	WB, IF, IHC-P, E
<b>Primary Accession</b>	<a href="#">P01286</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB14533
<b>Calculated MW</b>	12447
<b>Antigen Region</b>	8-34

## Additional Information

---

<b>Gene ID</b>	2691
<b>Other Names</b>	Somatoliberin, Growth hormone-releasing factor, GRF, Growth hormone-releasing hormone, GHRH, Somatocrinin, Somatorelin, Sermorelin, GHRH, GHRF
<b>Target/Specificity</b>	This GHRH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-34 amino acids from the N-terminal region of human GHRH.
<b>Dilution</b>	WB~~1:1000 IF~~1:10~50 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	GHRH Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GHRH
<b>Synonyms</b>	GHRF

<b>Function</b>	GRF is released by the hypothalamus and acts on the adenohypophyse to stimulate the secretion of growth hormone.
<b>Cellular Location</b>	Secreted.


## Background

---

GHRH belongs to the glucagon family and is a preproprotein that is produced in the hypothalamus. The preproprotein is cleaved to form a 44 aa factor, also called somatocrinin, that acts to stimulate growth hormone release from the pituitary. Variant receptors for somatocrinin have been found in several types of tumors, and antagonists of these receptors can inhibit the growth of the tumors. Defects in this protein are a cause of dwarfism, while hypersecretion of the encoded protein is a cause of gigantism.

## Images

---



All lanes : Anti-GHRH Antibody (N-term) at 1:500 dilution  
 Lane 1: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution.  
 Observed band size : 12kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.