

# Integrin beta 2 Rabbit mAb

Catalog # AP77667

## Product Information

---

<b>Application</b>	WB, IP
<b>Primary Accession</b>	<a href="#">P05107</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Immunogen</b>	A synthesized peptide derived from human CD18
<b>Purification</b>	Affinity Chromatography
<b>Calculated MW</b>	84791

## Additional Information

---

<b>Gene ID</b>	3689
<b>Other Names</b>	ITGB2
<b>Dilution</b>	WB~~1/500-1/1000 IP~~N/A
<b>Format</b>	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	ITGB2 ( <a href="#">HGNC:6155</a> )
<b>Synonyms</b>	CD18, MFI7
<b>Function</b>	Integrin ITGAL:ITGB2 is a receptor for ICAM1, ICAM2 and ICAM3 (PubMed: <a href="#">1676048</a> , PubMed: <a href="#">23775590</a> , PubMed: <a href="#">38195629</a> ). Integrin ITGAL:ITGB2 is also a receptor for the secreted form of ubiquitin-like protein ISG15; the interaction is mediated by ITGAL (PubMed: <a href="#">29100055</a> ). Integrins ITGAM:ITGB2 and ITGAX:ITGB2 are receptors for the iC3b fragment of the third complement component and for fibrinogen. Integrin ITGAX:ITGB2 recognizes the sequence G-P-R in fibrinogen alpha-chain. Integrin ITGAM:ITGB2 recognizes P1 and P2 peptides of fibrinogen gamma chain. Integrin ITGAM:ITGB2 is also a receptor for factor X. Integrin ITGAD:ITGB2 is a receptor for ICAM3 and VCAM1 (PubMed: <a href="#">10438935</a> , PubMed: <a href="#">8777714</a> , PubMed: <a href="#">9841932</a> ). Contributes to natural killer cell cytotoxicity (PubMed: <a href="#">15356110</a> ). Involved in leukocyte adhesion and transmigration of

leukocytes including T-cells and neutrophils (PubMed:[11812992](#), PubMed:[28807980](#)). Triggers neutrophil transmigration during lung injury through PTK2B/PYK2-mediated activation (PubMed:[18587400](#)). Integrin ITGAL:ITGB2 in association with ICAM3, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed:[23775590](#)). In association with alpha subunit ITGAM/CD11b, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:[21193407](#)). Integrins ITGAX:ITGB2 functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis (PubMed:[16985175](#)). Integrins ITGAX:ITGB2 functions as a receptor of the neuron-specific adhesion molecule ICAM5 ensuring neuron cell-leukocyte adhesion (PubMed:[10741396](#)). Integrin ITGAL:ITGB2 functions as a receptor of ICAM1 by acting as a platform at the immunological synapse to translate TCR engagement and density of the ITGAL ligand ICAM1 into graded adhesion (PubMed:[38195629](#)). Integrin ITGAM:ITGB2/MAC-1 complex functions as a signaling receptor for the ligand receptor ICAM1, ensuring adhesion between stimulated neutrophils and stimulated endothelial cells (PubMed:[1980124](#)). Integrin ITGAL/ITGB2 that functions as a signaling receptor of ICAM2, ensuring leukocyte cell-cell adhesion on resting cells (PubMed:[1676048](#)).

#### Cellular Location


Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein

#### Tissue Location

Leukocytes (PubMed:23775590). Expressed in neutrophils (at protein level) (PubMed:21193407, PubMed:28807980)

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.