

PRAS40 Rabbit mAb

Catalog # AP77699

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	Q96B36
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human PRAS40
Purification	Affinity Chromatography
Calculated MW	27383

Additional Information

Gene ID	84335
Other Names	AKT1S1
Dilution	WB~~1/500-1/1000 IHC-P~~N/A FC~~1:10~50 IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	AKT1S1 {ECO:0000312 EMBL:AAH16043.1}
Function	Negative regulator of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed: 17277771 , PubMed: 17386266 , PubMed: 17510057 , PubMed: 29236692). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1 activity by blocking the MTOR substrate- recruitment site (PubMed: 29236692). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed: 17386266 , PubMed: 18372248). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed: 16174443 , PubMed: 18372248). May also play a role in nerve growth factor-mediated neuroprotection (By similarity).
Cellular Location	Cytoplasm, cytosol {ECO:0000250 UniProtKB:Q9D1F4}. Note=Found in the

cytosolic fraction of the brain. {ECO:0000250|UniProtKB:Q9D1F4}

Tissue Location

Widely expressed with highest levels of expression in liver and heart.
Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.