

# Periostin Rabbit mAb

Catalog # AP77702

#### **Product Information**

**Application** WB, IF, FC, ICC

Primary Accession
Reactivity
Human
Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human Periostin

**Purification** Affinity Chromatography

Calculated MW 93314

### **Additional Information**

**Gene ID** 10631

Other Names POSTN

**Dilution** WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name POSTN

**Synonyms** OSF2

**Function** Induces cell attachment and spreading and plays a role in cell adhesion

(PubMed:<u>12235007</u>). Enhances incorporation of BMP1 in the fibronectin matrix of connective tissues, and subsequent proteolytic activation of lysyl

oxidase LOX (By similarity).

**Cellular Location** Golgi apparatus {ECO:0000250 | UniProtKB:Q62009}. Secreted Secreted,

extracellular space, extracellular matrix Note=Colocalizes with BMP1 in the


Golgi {ECO:0000250 | UniProtKB:Q62009}

**Tissue Location** Widely expressed with highest levels in aorta, stomach, lower gastrointestinal

tract, placenta, uterus, thyroid tissue and breast. Expressed in the kidney (PubMed:21763681). Expressed in the lung (PubMed:22079858). Up-regulated

in epithelial ovarian tumors. Not expressed in normal ovaries. Also highly expressed at the tumor periphery of lung carcinoma tissue but not within the tumor Overexpressed in breast cancers.

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.