

DHFR Rabbit mAb

Catalog # AP77704

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	P00374
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human DHFR
Purification	Affinity Chromatography
Calculated MW	21453

Additional Information

Gene ID	1719
Other Names	DHFR
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	DHFR (HGNC:2861)
Function	Catalyzes the reduction of 7,8-dihydrofolate (DHF) to 5,6,7,8-tetrahydrofolate in a NADPH-dependent manner (PubMed: 12096917 , PubMed: 15039552 , PubMed: 17569517 , PubMed: 19196009 , PubMed: 19478082 , PubMed: 21876184 , PubMed: 9719595). Key enzyme in folate metabolism. Contributes to the nuclear and mitochondrial de novo thymidylate biosynthesis pathway (PubMed: 21876188 , PubMed: 22235121). Catalyzes an essential reaction for de novo glycine and purine synthesis, and for DNA precursor synthesis. Binds its own mRNA and that of DHFR2.
Cellular Location	Mitochondrion {ECO:0000250 UniProtKB:P00375}. Cytoplasm {ECO:0000250 UniProtKB:P00375}. Nucleus. Note=Localized to the nucleus during S and G2/M phases of the cell cycle. As a component of the de novo thymidylate synthesis complex, localizes specifically to replication forks during DNA synthesis (PubMed:22235121)

Tissue Location

Widely expressed in fetal and adult tissues, including throughout the fetal and adult brains and whole blood Expression is higher in the adult brain than in the fetal brain

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.