

## CXCR3 Rabbit mAb

Catalog # AP77709

### **Product Information**

**Application** WB, IF, ICC, IP

Primary Accession P49682
Reactivity Human
Host Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human CXCR3

**Purification** Affinity Chromatography

Calculated MW 40660

### **Additional Information**

Gene ID 2833

Other Names CXCR3

**Dilution** WB~~1/500-1/1000 IF~~1:50~200 ICC~~N/A IP~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name CXCR3

**Synonyms** GPR9

**Function** [Isoform 1]: Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11

and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G- protein signaling pathway (PubMed:12782716). Binds to CCL21. Probably promotes cell chemotaxis response. Upon activation by PF4, induces activated T-lymphocytes migration mediated via downstream Ras/extracellular signal-regulated kinase (ERK)


signaling. [Isoform 3]: Mediates the activity of CXCL11.

**Cellular Location** [Isoform 1]: Cell membrane; Multi-pass membrane protein

**Tissue Location** Isoform 1 and isoform 2 are mainly expressed in heart, kidney, liver and

skeletal muscle. Isoform 1 is also expressed in placenta. Isoform 2 is

# **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.