

Cytochrome P450 3A4 Rabbit mAb

Catalog # AP77739

Product Information

Application	WB, IHC-P, IP
Primary Accession	P08684
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Cytochrome P450 3A4
Purification	Affinity Chromatography
Calculated MW	57343

Additional Information

Gene ID	1576
Other Names	CYP3A4
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CYP3A4 {ECO:0000303 PubMed:11470997, ECO:0000312 HGNC:HGNC:2637}
Function	A cytochrome P450 monooxygenase involved in the metabolism of sterols, steroid hormones, retinoids and fatty acids (PubMed:10681376, PubMed:11093772, PubMed:11555828, PubMed:12865317, PubMed:14559847, PubMed:15373842, PubMed:15764715, PubMed:19965576, PubMed:20702771, PubMed:21490593, PubMed:21576599). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase). Catalyzes the hydroxylation of carbon-hydrogen bonds (PubMed:12865317, PubMed:14559847, PubMed:15373842, PubMed:15764715, PubMed:21490593, PubMed:21576599, PubMed:2732228). Exhibits high catalytic activity for the formation of hydroxysteroids from estrone (E1) and 17beta- estradiol (E2),

namely 2-hydroxy E1 and E2, as well as D-ring hydroxylated E1 and E2 at the C-16 position (PubMed:[11555828](#), PubMed:[12865317](#), PubMed:[14559847](#)). Plays a role in the metabolism of androgens, particularly in oxidative deactivation of testosterone (PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[22773874](#), PubMed:[2732228](#)). Metabolizes testosterone to less biologically active 2beta- and 6beta- hydroxytestosterones (PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[2732228](#)). Contributes to the formation of hydroxycholesterols (oxysterols), particularly A-ring hydroxylated cholesterol at the C- 4beta position, and side chain hydroxylated cholesterol at the C-25 position, likely contributing to cholesterol degradation and bile acid biosynthesis (PubMed:[21576599](#)). Catalyzes bisallylic hydroxylation of polyunsaturated fatty acids (PUFA) (PubMed:[9435160](#)). Catalyzes the epoxidation of double bonds of PUFA with a preference for the last double bond (PubMed:[19965576](#)). Metabolizes endocannabinoid arachidonoyl ethanolamide (anandamide) to 8,9-, 11,12-, and 14,15- epoxyeicosatrienoic acid ethanolamides (EpETE-EAs), potentially modulating endocannabinoid system signaling (PubMed:[20702771](#)). Plays a role in the metabolism of retinoids. Displays high catalytic activity for oxidation of all-trans-retinol to all-trans-retinal, a rate- limiting step for the biosynthesis of all-trans-retinoic acid (atRA) (PubMed:[10681376](#)). Further metabolizes atRA toward 4-hydroxyretinoate and may play a role in hepatic atRA clearance (PubMed:[11093772](#)). Responsible for oxidative metabolism of xenobiotics. Acts as a 2-exo- monooxygenase for plant lipid 1,8-cineole (eucalyptol) (PubMed:[11159812](#)). Metabolizes the majority of the administered drugs. Catalyzes sulfoxidation of the anthelmintics albendazole and fenbendazole (PubMed:[10759686](#)). Hydroxylates antimalarial drug quinine (PubMed:[8968357](#)). Acts as a 1,4-cineole 2-exo-monooxygenase (PubMed:[11695850](#)). Also involved in vitamin D catabolism and calcium homeostasis. Catalyzes the inactivation of the active hormone calcitriol (1-alpha,25-dihydroxyvitamin D(3)) (PubMed:[29461981](#)).


Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein.
 Microsome membrane; Single-pass membrane protein

Tissue Location

Expressed in prostate and liver. According to some authors, it is not expressed in brain (PubMed:[19094056](#)). According to others, weak levels of expression are measured in some brain locations (PubMed:[18545703](#), PubMed:[19359404](#)). Also expressed in epithelium of the small intestine and large intestine, bile duct, nasal mucosa, kidney, adrenal cortex, epithelium of the gastric mucosa with intestinal metaplasia, gallbladder, intercalated ducts of the pancreas, chief cells of the parathyroid and the corpus luteum of the ovary (at protein level).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.