10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Natriuretic Peptides A Rabbit mAb

Catalog # AP77803

Product Information

Application	WB, IP
Primary Accession	P01160
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human ANP
Purification	Affinity Chromatography
Calculated MW	16396

Additional Information

Gene ID	4878
Other Names	NPPA
Dilution	WB~~1/500-1/1000 IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	NPPA
Synonyms	ANP, PND
Function	[Atrial natriuretic peptide]: Hormone that plays a key role in mediating cardio-renal homeostasis, and is involved in vascular remodeling and regulating energy metabolism (PubMed:15741263, PubMed:16875975, PubMed:18835931, PubMed:21672517, PubMed:22307324, PubMed:2532366, PubMed:2825692, PubMed:7595132, PubMed:7720651, PubMed:8087923, PubMed:8653797). Acts by specifically binding and stimulating NPR1 to produce cGMP, which in turn activates effector proteins, such as PRKG1, that drive various biological responses (PubMed:1660465, PubMed:1672777, PubMed:21098034, PubMed:2162527, PubMed:22307324, PubMed:2825692, PubMed:7720651, PubMed:8384600, PubMed:9893117). Regulates vasodilation, natriuresis, diuresis and aldosterone synthesis and is therefore essential for regulating blood pressure, controlling the extracellular fluid

volume and maintaining the fluid-electrolyte balance (PubMed:2532366, PubMed:2825692, PubMed:7595132, PubMed:7720651, PubMed:8087923, PubMed:8653797). Also involved in inhibiting cardiac remodeling and cardiac hypertrophy by inducing cardiomyocyte apoptosis and attenuating the growth of cardiomyocytes and fibroblasts (PubMed:16875975). Plays a role in female pregnancy by promoting trophoblast invasion and spiral artery remodeling in uterus, and thus prevents pregnancy-induced hypertension (By similarity). In adipose tissue, acts in various cGMP- and PKG- dependent pathways to regulate lipid metabolism and energy homeostasis (PubMed:15741263, PubMed:18835931, PubMed:21672517, PubMed:22307324). This includes up-regulating lipid metabolism and mitochondrial oxygen utilization by activating the AMP-activated protein kinase (AMPK), and increasing energy expenditure by acting via MAPK11 to promote the UCP1- dependent thermogenesis of brown adipose tissue (PubMed:15741263, PubMed:18835931, PubMed:21672517, PubMed:22307324). Binds the clearance receptor NPR3 which removes the hormone from circulation (PubMed:1672777).


Cellular Location

[Long-acting natriuretic peptide]: Secreted. Note=Detected in blood. [Kaliuretic peptide]: Secreted. Note=Detected in blood [Atrial natriuretic peptide]: Secreted. Perikaryon. Cell projection. Note=Detected in blood (PubMed:15741263, PubMed:18835931, PubMed:2532366, PubMed:7955907, PubMed:7984506, PubMed:8351194, PubMed:8653797, PubMed:8779891). Detected in urine in one study (PubMed:8351194). However, in another study, was not detected in urine (PubMed:7984506). Detected in cytoplasmic bodies and neuronal processes of pyramidal neurons (layers II-VI) (PubMed:30534047) Increased secretion in response to the vasopressin AVP (By similarity) Likely to be secreted in response to an increase in atrial pressure or atrial stretch (PubMed:2532366). In kidney cells, secretion increases in response to activated guanylyl cyclases and increased intracellular cAMP levels (PubMed:9893117). Plasma levels increase 15 minutes after a high-salt meal, and decrease back to normal plasma levels 1 hr later (PubMed:8779891). {ECO:0000250|UniProtKB:P01161, ECO:0000269|PubMed:15741263, ECO:0000269|PubMed:18835931, ECO:0000269|PubMed:2532366, ECO:0000269|PubMed:30534047, ECO:0000269|PubMed:7955907, ECO:0000269|PubMed:7984506, ECO:0000269|PubMed:8351194, ECO:0000269|PubMed:8653797, ECO:0000269|PubMed:8779891, ECO:0000269|PubMed:9893117}

Tissue Location

[Urodilatin]: Detected in the kidney distal tubular cells (at protein level) (PubMed:8384600, PubMed:9794555). Present in urine (at protein level) (PubMed:2972874, PubMed:8351194, PubMed:8779891, PubMed:9794555).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.