10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MVP Rabbit mAb

Catalog # AP77816

Product Information

Application	WB, IHC-P
Primary Accession	Q14764
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human MVP
Purification	Affinity Chromatography
Calculated MW	99327

Additional Information

Gene ID	9961
Other Names	MVP
Dilution	WB~~1/500-1/1000 IHC-P~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	MVP
Synonyms	LRP
Function	Structural component of the vault particle, which is a massive ribonucleoprotein complex that may play a role in cargo transport and signal transduction (PubMed:40691181). Oligomerizes to form a cage-like vault complex, which can house several minor vault components including PARP4 (PubMed:40691181). Vaults may play a role in nucleo-cytoplasmic transport (PubMed:15133037). Down-regulates IFNG- mediated STAT1 signaling and subsequent activation of JAK (PubMed:16418217). Down-regulates SRC activity and signaling through MAP kinases (PubMed:16441665).
Cellular Location	Cytoplasm. Nucleus, nuclear pore complex. Cytoplasm, perinuclear region. Note=5% found in the nuclear pore complex (PubMed:15133037). Translocates from the nucleus to the cytoplasm upon EGF treatment

(PubMed:16441665)

Tissue Location

Present in most normal tissues. Higher expression observed in epithelial cells with secretory and excretory functions, as well as in cells chronically exposed to xenobiotics, such as bronchial cells and cells lining the intestine. Overexpressed in many multidrug- resistant cancer cells

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.