

# HBEGF Rabbit mAb

Catalog # AP77969

## Product Information

---

|                          |                                                 |
|--------------------------|-------------------------------------------------|
| <b>Application</b>       | WB                                              |
| <b>Primary Accession</b> | <a href="#">Q99075</a>                          |
| <b>Reactivity</b>        | Rat, Human                                      |
| <b>Host</b>              | Rabbit                                          |
| <b>Clonality</b>         | Monoclonal Antibody                             |
| <b>Isotype</b>           | IgG                                             |
| <b>Conjugate</b>         | Unconjugated                                    |
| <b>Immunogen</b>         | A synthesized peptide derived from human HB EGF |
| <b>Purification</b>      | Affinity Chromatography                         |
| <b>Calculated MW</b>     | 23067                                           |

## Additional Information

---

|                    |                                                                                                    |
|--------------------|----------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>     | 1839                                                                                               |
| <b>Other Names</b> | HBEGF                                                                                              |
| <b>Dilution</b>    | WB~~1/500-1/1000                                                                                   |
| <b>Format</b>      | Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol. |
| <b>Storage</b>     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.           |


## Protein Information

---

|                          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Name</b>              | HBEGF                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Synonyms</b>          | DTR, DTS, HEGFL                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Function</b>          | Growth factor that mediates its effects via EGFR, ERBB2 and ERBB4. Required for normal cardiac valve formation and normal heart function. Promotes smooth muscle cell proliferation. May be involved in macrophage-mediated cellular proliferation. It is mitogenic for fibroblasts, but not endothelial cells. It is able to bind EGF receptor/EGFR with higher affinity than EGF itself and is a far more potent mitogen for smooth muscle cells than EGF. Also acts as a diphtheria toxin receptor. |
| <b>Cellular Location</b> | [Heparin-binding EGF-like growth factor]: Secreted, extracellular space. Note=Mature HB-EGF is released into the extracellular space and probably binds to a receptor                                                                                                                                                                                                                                                                                                                                  |

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.