

# APOC3 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7797b

## Product Information

---

<b>Application</b>	IHC-P, WB, E
<b>Primary Accession</b>	<a href="#">P02656</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB15169
<b>Calculated MW</b>	10852
<b>Antigen Region</b>	70-99

## Additional Information

---

<b>Gene ID</b>	345
<b>Other Names</b>	Apolipoprotein C-III, Apo-CIII, ApoC-III, Apolipoprotein C3, APOC3
<b>Target/Specificity</b>	This APOC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 70~99 amino acids from the C-term region of human APOC3.
<b>Dilution</b>	IHC-P~~1:100~500 WB~~1:2000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	APOC3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	APOC3
<b>Function</b>	Component of triglyceride-rich very low density lipoproteins (VLDL) and high density lipoproteins (HDL) in plasma (PubMed: <a href="#">18201179</a> , PubMed: <a href="#">22510806</a> ). Plays a multifaceted role in triglyceride homeostasis (PubMed: <a href="#">18201179</a> , PubMed: <a href="#">22510806</a> ). Intracellularly, promotes hepatic very low density lipoprotein 1 (VLDL1) assembly and secretion; extracellularly, attenuates

hydrolysis and clearance of triglyceride- rich lipoproteins (TRLs) (PubMed:[18201179](#), PubMed:[22510806](#)). Impairs the lipolysis of TRLs by inhibiting lipoprotein lipase and the hepatic uptake of TRLs by remnant receptors (PubMed:[18201179](#), PubMed:[22510806](#)). Formed of several curved helices connected via semiflexible hinges, so that it can wrap tightly around the curved micelle surface and easily adapt to the different diameters of its natural binding partners (PubMed:[18408013](#)).

**Cellular Location** Secreted

**Tissue Location** Liver..

## Background

---

Apolipoprotein C-III is a very low density lipoprotein (VLDL) protein. APOC3 inhibits lipoprotein lipase and hepatic lipase; it is thought to delay catabolism of triglyceride-rich particles. The APOA1, APOC3 and APOA4 genes are closely linked in both rat and human genomes. The A-I and A-IV genes are transcribed from the same strand, while the A-1 and C-III genes are convergently transcribed. An increase in apoC-III levels induces the development of hypertriglyceridemia.

## References


---

Liu,Y., Pharmacogenet. Genomics 19 (2), 161-169 (2009)

Pollin,T.I., Science 322 (5908), 1702-1705 (2008)

## Images

---



Anti-APOC3 Antibody (C-term) at 1:2000 dilution + human plasma lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 11 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.