10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Rab25 Rabbit mAb

Catalog # AP78008

Product Information

Application	WB
Primary Accession	P57735
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Rab25
Purification	Affinity Chromatography
Calculated MW	23496

Additional Information

Gene ID	57111
Other Names	RAB25
Dilution	WB~~1/500-1/1000
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	RAB25 (HGNC:18238)
Synonyms	CATX8
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (By similarity). RAB25 regulates epithelial cell differentiation, proliferation and survival, thereby playing key roles in tumorigenesis (PubMed:17925226). Promotes invasive migration of cells in which it functions to localize and maintain integrin alpha-V/beta-1 at the tips of extending pseudopodia (PubMed:17925226). Involved in the regulation of epithelial morphogenesis through the control of CLDN4 expression and localization at tight junctions (By similarity). May selectively regulate the

	apical recycling pathway (By similarity). Together with MYO5B regulates transcytosis (By similarity).
Cellular Location	Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic vesicle. Cell projection, pseudopodium membrane. Note=Colocalizes with integrin alpha-V/beta-1 in vesicles at the pseudopodial tips. Colocalizes with RAB11A in subapical vesicles (By similarity). {ECO:0000250|UniProtKB:P46629, ECO:0000269|PubMed:17925226}
Tissue Location	Expression is restricted to epithelial cells (PubMed:15502842). Expressed in ovarian epithelium (NOE) and breast tissue. Expressed in ovarian cancer; expression is increased relative to NOE cells. Expression in ovarian cancer is stage dependent, with stage III and stage IV showing higher levels than early stage cancers Expressed in breast cancer; expression is increased relative to normal breast tissue.

Images

Image not found : 202310/R383076-WB-1.jpg

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.