

MARK3 Rabbit mAb

Catalog # AP78123

Product Information

Application	WB, IF, FC, ICC, IP
Primary Accession	P27448
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human MARK3
Purification	Affinity Chromatography
Calculated MW	84429

Additional Information

Gene ID	4140
Other Names	MARK3
Dilution	WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	MARK3
Synonyms	CTAK1, EMK2
Function	Serine/threonine-protein kinase (PubMed: 16822840 , PubMed: 16980613 , PubMed: 23666762). Involved in the specific phosphorylation of microtubule-associated proteins for MAP2 and MAP4. Phosphorylates the microtubule-associated protein MAPT/TAU (PubMed: 23666762). Phosphorylates CDC25C on 'Ser-216' (PubMed: 12941695). Regulates localization and activity of some histone deacetylases by mediating phosphorylation of HDAC7, promoting subsequent interaction between HDAC7 and 14-3-3 and export from the nucleus (PubMed: 16980613). Regulates localization and activity of MITF by mediating its phosphorylation, promoting subsequent interaction between MITF and 14-3-3 and retention in the cytosol (PubMed: 16822840). Negatively regulates the Hippo signaling pathway and antagonizes the phosphorylation of LATS1. Cooperates with

DLG5 to inhibit the kinase activity of STK3/MST2 toward LATS1 (PubMed:[28087714](#)). Phosphorylates PKP2 and KSR1 (PubMed:[12941695](#)).


Cellular Location

Cell membrane; Peripheral membrane protein. Cell projection, dendrite. Cytoplasm

Tissue Location

Ubiquitous.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.