

TPPP Rabbit mAb

Catalog # AP78138

Product Information

Application	WB, IHC-P, IF, FC, ICC
Primary Accession	O94811
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human TPPP
Purification	Affinity Chromatography
Calculated MW	23694

Additional Information

Gene ID	11076
Other Names	TPPP
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	TPPP {ECO:0000303 PubMed:17105200, ECO:0000312 HGNC:HGNC:24164}
Function	Regulator of microtubule dynamics that plays a key role in myelination by promoting elongation of the myelin sheath (PubMed: 31522887). Acts as a microtubule nucleation factor in oligodendrocytes: specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, and promotes microtubule nucleation, an important step for elongation of the myelin sheath (PubMed: 31522887 , PubMed: 33831707). Required for both uniform polarized growth of distal microtubules as well as directing the branching of proximal processes (PubMed: 31522887). Shows magnesium-dependent GTPase activity; the role of the GTPase activity is unclear (PubMed: 21316364 , PubMed: 21995432). In addition to microtubule nucleation activity, also involved in microtubule bundling and stabilization of existing microtubules, thereby maintaining the integrity of the microtubule network (PubMed: 17105200 , PubMed: 17693641 , PubMed: 18028908 , PubMed: 26289831). Regulates microtubule dynamics by promoting tubulin

acetylation: acts by inhibiting the tubulin deacetylase activity of HDAC6 (PubMed:[20308065](#), PubMed:[23093407](#)). Also regulates cell migration: phosphorylation by ROCK1 inhibits interaction with HDAC6, resulting in decreased acetylation of tubulin and increased cell motility (PubMed:[23093407](#)). Plays a role in cell proliferation by regulating the G1/S-phase transition (PubMed:[23355470](#)). Involved in astral microtubule organization and mitotic spindle orientation during early stage of mitosis; this process is regulated by phosphorylation by LIMK2 (PubMed:[22328514](#)).


Cellular Location

Golgi outpost {ECO:0000250|UniProtKB:D3ZQL7}. Cytoplasm, cytoskeleton, microtubule organizing center {ECO:0000250|UniProtKB:D3ZQL7}. Cytoplasm, cytoskeleton. Nucleus Cytoplasm, cytoskeleton, spindle Note=Specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, which shapes dendrite morphology by functioning as sites of acentrosomal microtubule nucleation (By similarity). Mainly localizes to the cytoskeleton (PubMed:18028908) Also found in the nucleus; however, nuclear localization is unclear and requires additional evidences (PubMed:18028908). Localizes to glial Lewy bodies in the brains of individuals with synucleinopathies (PubMed:15590652, PubMed:17027006). During mitosis, colocalizes with LIMK2 at the mitotic spindle (PubMed:22328514) {ECO:0000250|UniProtKB:D3ZQL7, ECO:0000269|PubMed:15590652, ECO:0000269|PubMed:17027006, ECO:0000269|PubMed:18028908, ECO:0000269|PubMed:22328514}

Tissue Location

Widely expressed..

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.