

## **ACADM Rabbit mAb**

Catalog # AP78154

#### **Product Information**

**Application** WB, IHC-P, IF, ICC, IP

Primary Accession P11310

Reactivity Rat, Human, Mouse

**Host** Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human ACADM

**Purification** Affinity Chromatography

Calculated MW 46588

### **Additional Information**

Gene ID 34

Other Names ACADM

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A IP~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name ACADM ( HGNC:89)

**Function** Medium-chain specific acyl-CoA dehydrogenase is one of the acyl-CoA

dehydrogenases that catalyze the first step of mitochondrial fatty acid beta-oxidation, an aerobic process breaking down fatty acids into acetyl-CoA


and allowing the production of energy from fats (PubMed:<u>1970566</u>, PubMed:<u>21237683</u>, PubMed:<u>2251268</u>, PubMed:<u>8823175</u>). The first step of

fatty acid beta-oxidation consists in the removal of one hydrogen from C-2 and C-3 of the straight-chain fatty acyl-CoA thioester, resulting in the formation of trans-2-enoyl-CoA (PubMed:2251268). Electron transfer

flavoprotein (ETF) is the electron acceptor that transfers electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase) (PubMed:15159392, PubMed:25416781). Among the different mitochondrial acyl-CoA dehydrogenases, medium-chain specific acyl-CoA dehydrogenase acts specifically on acyl-CoAs with saturated 6 to 12 carbons long primary chains (PubMed:1970566, PubMed:21237683, PubMed:2251268,

Mitochondrion matrix

# **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.