

PSMD14 Rabbit mAb

Catalog # AP78163

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	O00487
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human PSMD14
Purification	Affinity Chromatography
Calculated MW	34577

Additional Information

Gene ID	10213
Other Names	PSMD14
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	PSMD14
Synonyms	POH1
Function	Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair (PubMed: 9374539 , PubMed: 1317798). The PSMD14 subunit is a metalloprotease that specifically cleaves 'Lys-63'-linked polyubiquitin chains within the complex (PubMed: 22909820). Plays a role in response to double-strand breaks (DSBs): acts as a regulator of non-homologous end joining (NHEJ) by cleaving 'Lys-63'-linked polyubiquitin, thereby promoting retention of JMJD2A/KDM4A on chromatin and restricting

TP53BP1 accumulation (PubMed:[22909820](#)). Also involved in homologous recombination repair by promoting RAD51 loading (PubMed:[22909820](#)). Regulates macroautophagy by ensuring Golgi-to-ER retrograde transport through its deubiquitinating activity on K63-linked ubiquitin chains. This activity prevents the retention of essential autophagy proteins at the Golgi, enabling their trafficking to autophagosome formation sites and supporting Golgi-ER membrane recycling critical for effective autophagy (PubMed:[32210007](#)).

Tissue Location

Widely expressed. Highest levels in heart and skeletal muscle.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.