

## NRG3 Rabbit mAb

Catalog # AP78176

### **Product Information**

**Application** WB, FC **Primary Accession** P56975

**Reactivity** Rat, Human, Mouse

**Host** Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human NRG3

**Purification** Affinity Chromatography

Calculated MW 77901

#### **Additional Information**

**Gene ID** 10718

Other Names NRG3

**Dilution** WB~~1/500-1/1000 FC~~1:10~50

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name NRG3

**Function** Direct ligand for the ERBB4 tyrosine kinase receptor. Binding results in

ligand-stimulated tyrosine phosphorylation and activation of the receptor. Does not bind to the EGF receptor, ERBB2 or ERBB3 receptors. May be a

survival factor for oligodendrocytes.

**Cellular Location** [Pro-neuregulin-3, membrane-bound isoform]: Cell membrane; Single-pass

type I membrane protein. Note=Does not seem to be active. [Isoform 3]: Cell membrane; Single-pass type I membrane protein. Note=Isoform 3 is also


proteolytically released as a soluble form

**Tissue Location** Highly expressed in most regions of the brain with the exception of corpus

callosum. Expressed at lower level in testis Not detected in heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, ovary,

small intestine, colon and peripheral blood leukocytes

# **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.