

# EHMT2 Rabbit mAb

Catalog # AP78208

#### **Product Information**

Application WB, IHC-P
Primary Accession Q96KQ7
Reactivity Human
Host Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human EHMT2/G9A

**Purification** Affinity Purified

Calculated MW 132370

### **Additional Information**

**Gene ID** 10919

Other Names EHMT2

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name EHMT2

**Synonyms** BAT8, C6orf30, G9A, KMT1C, NG36

**Function** Histone methyltransferase that specifically mono- and dimethylates 'Lys-9'

of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by

recruiting HP1 proteins to methylated histones. Also mediates

monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In

addition to the histone methyltransferase activity, also methylates

non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

**Cellular Location** Nucleus. Chromosome. Note=Associates with euchromatic regions

(PubMed:11316813). Does not associate with heterochromatin


(PubMed:11316813).

**Tissue Location** Expressed in all tissues examined, with high levels in fetal liver, thymus,

lymph node, spleen and peripheral blood leukocytes and lower level in bone

marrow

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.