

# NEK6 Rabbit mAb

Catalog # AP78217

## Product Information

---

<b>Application</b>	WB, IP
<b>Primary Accession</b>	<a href="#">Q9HC98</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Immunogen</b>	A synthesized peptide derived from human NEK6
<b>Purification</b>	Affinity Purified
<b>Calculated MW</b>	35714

## Additional Information

---

<b>Gene ID</b>	10783
<b>Other Names</b>	NEK6
<b>Dilution</b>	WB~~1/500-1/1000 IP~~N/A
<b>Format</b>	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	NEK6 ( <a href="#">HGNC:7749</a> )
<b>Function</b>	Protein kinase which plays an important role in mitotic cell cycle progression (PubMed: <a href="#">11516946</a> , PubMed: <a href="#">14563848</a> ). Required for chromosome segregation at metaphase-anaphase transition, robust mitotic spindle formation and cytokinesis (PubMed: <a href="#">19414596</a> ). Phosphorylates ATF4, CIRSR, PTN, RAD26L, RBBP6, RPS7, RPS6KB1, TRIP4, STAT3 and histones H1 and H3 (PubMed: <a href="#">12054534</a> , PubMed: <a href="#">20873783</a> ). Phosphorylates KIF11 to promote mitotic spindle formation (PubMed: <a href="#">19001501</a> ). Involved in G2/M phase cell cycle arrest induced by DNA damage (PubMed: <a href="#">18728393</a> ). Inhibition of activity results in apoptosis. May contribute to tumorigenesis by suppressing p53/TP53-induced cancer cell senescence (PubMed: <a href="#">21099361</a> ). Phosphorylates EML4 at 'Ser-144', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed: <a href="#">31409757</a> ).

## Cellular Location


Cytoplasm. Nucleus. Nucleus speckle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole.  
Note=Colocalizes with APBB1 at the nuclear speckles. Colocalizes with PIN1 in the nucleus. Colocalizes with ATF4, CIRS1, ARHGAP33, ANKRA2, CDC42, NEK9, RAD26L, RBBP6, RPS7, TRIP4, RELB and PHF1 in the centrosome. Localizes to spindle microtubules in metaphase and anaphase and to the midbody during cytokinesis

## Tissue Location

Ubiquitous, with highest expression in heart and skeletal muscle.

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.