

# VPS11 Rabbit mAb

Catalog # AP78237

## Product Information

---

<b>Application</b>	WB, IHC-P, FC
<b>Primary Accession</b>	<a href="#">Q9H270</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Immunogen</b>	A synthesized peptide derived from human VPS11
<b>Purification</b>	Affinity Purified
<b>Calculated MW</b>	107837

## Additional Information

---

<b>Gene ID</b>	55823
<b>Other Names</b>	VPS11
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A FC~~1:10~50
<b>Format</b>	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	VPS11
<b>Synonyms</b>	RNF108
<b>Function</b>	Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Believed to act as a core component of the putative HOPS and CORVET endosomal tethering complexes which are proposed to be involved in the Rab5-to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes. The CORVET complex is proposed to function as a Rab5 effector to mediate early endosome fusion probably in specific endosome subpopulations (PubMed: <a href="#">11382755</a> , PubMed: <a href="#">23351085</a> , PubMed: <a href="#">24554770</a> ,

PubMed:[25266290](#), PubMed:[25783203](#)). Required for fusion of endosomes and autophagosomes with lysosomes (PubMed:[25783203](#)). Involved in cargo transport from early to late endosomes and required for the transition from early to late endosomes (PubMed:[21148287](#)). Involved in the retrograde Shiga toxin transport (PubMed:[23593995](#)).

#### Cellular Location


Endosome. Late endosome membrane; Peripheral membrane protein; Cytoplasmic side. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome {ECO:0000269 | PubMed:21148287, ECO:0000305}. Cytoplasmic vesicle. Cytoplasmic vesicle, autophagosome. Cytoplasmic vesicle, clathrin-coated vesicle

#### Tissue Location

Ubiquitous. Expression was highest in heart and low in lung

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.