

DcR2 Rabbit mAb

Catalog # AP78306

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	Q9UBN6
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human DCR2
Purification	Affinity Purified
Calculated MW	41849


Additional Information

Gene ID	8793
Other Names	TNFRSF10D
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	TNFRSF10D (HGNC:11907)
Function	Receptor for the cytotoxic ligand TRAIL (PubMed: 9430226). Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis (PubMed: 9537512). Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to PubMed: 9382840 , it cannot but according to PubMed: 9430226 , it can induce the NF-kappa-B pathway (PubMed: 9382840 , PubMed: 9430226).
Cellular Location	Membrane; Single-pass type I membrane protein
Tissue Location	Widely expressed, in particular in fetal kidney, lung and liver, and in adult testis and liver. Also expressed in peripheral blood leukocytes, colon and small intestine, ovary, prostate, thymus, spleen, pancreas, kidney, lung, placenta and heart

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.