

Muscarinic Acetylcholine Receptor M2 Rabbit mAb

Catalog # AP78318

Product Information

Application	WB, IHC-P, IP
Primary Accession	P08172
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human ACM2
Purification	Affinity Purified
Calculated MW	51715


Additional Information

Gene ID	1129
Other Names	CHRM2
Dilution	WB~1/500-1/1000 IHC-P~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CHRM2
Function	Muscarinic receptor for acetylcholine, a neurotransmitter found in the brain, neuromuscular junctions and the autonomic ganglia (PubMed: 24256733 , PubMed: 3443095 , PubMed: 36690613). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (PubMed: 36690613). CHRM2 is coupled to G(i)/G(o) (GNAI1 or GNAO1) G proteins and mediates signaling by inhibiting adenylate cyclase activity (PubMed: 36690613).
Cellular Location	Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane protein. Note=Phosphorylation in response to agonist binding promotes receptor internalization {ECO:0000250 UniProtKB:P06199}

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.