

Lipoamide Dehydrogenase Rabbit mAb

Catalog # AP78392

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	P09622
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human DLDH
Purification	Affinity Purified
Calculated MW	54177

Additional Information

Gene ID	1738
Other Names	DLD
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	DLD (HGNC:2898)
Synonyms	GCSL, LAD, PHE3
Function	Lipoamide dehydrogenase is a component of the glycine cleavage system as well as an E3 component of three alpha-ketoacid dehydrogenase complexes (pyruvate-, alpha-ketoglutarate-, and branched- chain amino acid-dehydrogenase complex) (PubMed: 15712224 , PubMed: 16442803 , PubMed: 16770810 , PubMed: 17404228 , PubMed: 20160912 , PubMed: 20385101). The 2-oxoglutarate dehydrogenase complex is mainly active in the mitochondrion (PubMed: 29211711). A fraction of the 2-oxoglutarate dehydrogenase complex also localizes in the nucleus and is required for lysine succinylation of histones: associates with KAT2A on chromatin and provides succinyl-CoA to histone succinyltransferase KAT2A (PubMed: 29211711). In monomeric form may have additional moonlighting function as serine protease (PubMed: 17404228). Involved in the

hyperactivation of spermatazoa during capacitation and in the spermatazoal acrosome reaction (By similarity). The pyruvate dehydrogenase (PDH) complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2), and thereby links cytoplasmic glycolysis and the mitochondrial tricarboxylic acid (TCA) cycle (Probable). It contains multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and dihydrolipoamide dehydrogenase (E3) (Probable). The E3 subunit catalyzes reoxidation of the dihydrolipoyl moiety on lipoyl-bearing domains (LBDs) of E2 with NAD⁺ as the ultimate electron acceptor (PubMed:[16442803](#), PubMed:[16770810](#), PubMed:[20160912](#), PubMed:[20385101](#)).

Cellular Location

Mitochondrion matrix. Nucleus. Cell projection, cilium, flagellum {ECO:0000250|UniProtKB:Q811C4}. Cytoplasmic vesicle, secretory vesicle, acrosome. Note=Mainly localizes in the mitochondrion. A small fraction localizes to the nucleus, where the 2-oxoglutarate dehydrogenase complex is required for histone succinylation.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.