

ADK Rabbit mAb

Catalog # AP78471

Product Information

Primary Accession	P55263
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	40545


Additional Information

Gene ID	132
Other Names	ADK
Format	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ADK {ECO:0000303 PubMed:19635462, ECO:0000312 HGNC:HGNC:257}
Function	Adenosine kinase that mediates the phosphorylation of the purine nucleoside adenosine at the 5' position in an ATP-dependent manner: catalyzes phosphorylation of both unmodified and modified adenosines (PubMed:21963049, PubMed:40840445, PubMed:6246102, PubMed:8577746, PubMed:9070863). Plays a key role in the detoxification of modified adenosines containing N(6)-methylated adenine (m6A) post-transcriptional modification (PubMed:40840445). Modified nucleosides are derived from the degradation of RNAs (mRNAs, rRNAs and tRNAs) and possess intrinsic cytotoxicity and must be cleared to prevent metabolic dysfunction (PubMed:40840445). Catalyzes the phosphorylation of the free cytosolic methylated adenosine nucleotides N(6)-methyladenosine (m6A), N(6),N(6)-dimethyladenosine (m6,6A) and N(6)-isopentenyladenosine (i6A) into adenosine monophosphate (AMP) intermediates that are further detoxified by MAPDA/ADAL (PubMed:40840445).
Cellular Location	Cytoplasm, cytosol. [Isoform 2]: Cytoplasm
Tissue Location	Widely expressed. Highest level in placenta, liver, muscle and kidney.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.