

SLC27A4 Rabbit mAb

Catalog # AP78474

Product Information

Application WB, IF, FC, ICC **Primary Accession** Q6P1M0

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human SLC27A4 / FATP4

Purification Affinity Purified

Calculated MW 72064

Additional Information

Gene ID 10999

Other Names SLC27A4

Dilution WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name SLC27A4 (<u>HGNC:10998</u>)

Function Mediates the levels of long-chain fatty acids (LCFA) in the cell by facilitating

their transport across cell membranes (PubMed: 10518211, PubMed: 12556534, PubMed: 2048275, PubMed: 21395585

PubMed:<u>12556534</u>, PubMed:<u>20448275</u>, PubMed:<u>21395585</u>,

PubMed:<u>22022213</u>). Appears to be the principal fatty acid transporter in small intestinal enterocytes (PubMed:<u>20448275</u>). Also functions as an acyl-CoA ligase catalyzing the ATP-dependent formation of fatty acyl- CoA using LCFA and very-long-chain fatty acids (VLCFA) as substrates, which prevents fatty

acid efflux from cells and might drive more fatty acid uptake

(PubMed:<u>22022213</u>, PubMed:<u>24269233</u>). Plays a role in the formation of the epidermal barrier. Required for fat absorption in early embryogenesis (By similarity). Probably involved in fatty acid transport across the blood barrier (PubMed:<u>21395585</u>). Indirectly inhibits RPE65 via substrate competition and

via production of VLCFA derivatives like lignoceroyl-CoA. Prevents light-induced degeneration of rods and cones (By similarity).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Expressed at highest levels in brain, testis, colon and kidney. Expressed at medium levels in heart and liver, small intestine and stomach. Expressed at low levels in peripheral leukocytes, bone marrow, skeletal muscle and aorta. Expressed in adipose tissue (PubMed:24269233, PubMed:9878842). Expressed in brain gray matter (PubMed:21395585).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.