10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



LRP6 Rabbit mAb

Catalog # AP78525

Product Information

Application WB, FC, ICC **Primary Accession** 075581

Reactivity Human, Mouse, Rat

Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 180429

Additional Information

Gene ID 4040

Other Names LRP6

Dilution WB~~1/500-1/1000 FC~~1:10~50 ICC~~N/A

Format 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide

and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name LRP6

Function Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin

signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes (PubMed:11357136, PubMed:11448771,

PubMed: 15778503, PubMed: 16341017, PubMed: 16513652,

PubMed: 17326769, PubMed: 17400545, PubMed: 19107203,

PubMed:<u>19293931</u>, PubMed:<u>19801552</u>, PubMed:<u>28341812</u>). Cell-surface coreceptor of Wnt/beta- catenin signaling, which plays a pivotal role in bone formation (PubMed:<u>11357136</u>, PubMed:<u>11448771</u>, PubMed:<u>15778503</u>,

PubMed:<u>16341017</u>, PubMed:<u>16513652</u>, PubMed:<u>17326769</u>, PubMed:<u>17400545</u>, PubMed:<u>19107203</u>, PubMed:<u>19293931</u>,

PubMed: 19801552, PubMed: 28341812). The Wnt-induced Fzd/LRP6

coreceptor complex recruits DVL1 polymers to the plasma membrane which, in turn, recruits the AXIN1/GSK3B-complex to the cell surface promoting the

formation of signalosomes and inhibiting AXIN1/GSK3-mediated phosphorylation and destruction of beta-catenin (PubMed: 16513652). Required for posterior patterning of the epiblast during gastrulation (By

similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein Endoplasmic reticulum. Membrane raft. Note=On Wnt signaling, undergoes a cycle of caveolin- or clathrin-mediated endocytosis and plasma membrane location. Released from the endoplasmic reticulum on palmitoylation Mono-ubiquitination retains it in the endoplasmic reticulum in the absence of palmitoylation. On Wnt signaling, phosphorylated, aggregates and colocalizes with AXIN1 and GSK3B at the plasma membrane in LRP6- signalosomes (By similarity). Chaperoned to the plasma membrane by HSP90B1 and MESD (PubMed:23572575). {ECO:0000250 | UniProtKB:O88572, ECO:0000269 | PubMed:23572575}

Tissue Location

Widely coexpressed with LRP5 during embryogenesis and in adult tissues

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.