

# Phospho-EGFR (Tyr1086) Rabbit mAb

Catalog # AP78550

## **Product Information**

Application WB, IF, ICC, IP
Primary Accession P00533
Reactivity Human
Host Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human Phospho-EGFR (Y1086)

**Purification** Affinity Chromatography

Calculated MW 134277

# **Additional Information**

**Gene ID** 1956

Other Names EGFR

**Dilution** WB~~1/500-1/1000 IF~~1:50~200 ICC~~N/A IP~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

### **Protein Information**

Name EGFR ( HGNC:3236)

**Synonyms** ERBB, ERBB1, HER1

**Function** Receptor tyrosine kinase binding ligands of the EGF family and activating

several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:<u>10805725</u>, PubMed:<u>27153536</u>, PubMed:<u>2790960</u>, PubMed:<u>35538033</u>). Known ligands include EGF, TGFA/TGF- alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding

EGF (PubMed: 12297049, PubMed: 15611079, PubMed: 17909029,

PubMed: 20837704, PubMed: 27153536, PubMed: 2790960, PubMed: 7679104, PubMed: 8144591, PubMed: 9419975). Ligand binding triggers receptor homoand/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK,

PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity).


#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome. Endosome membrane. Nucleus. Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:17909029, PubMed:20674546). Endocytosed upon activation by ligand (PubMed:17182860, PubMed:17909029, PubMed:27153536, PubMed:2790960). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055)

#### **Tissue Location**

Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.