

DPF2 Rabbit mAb

Catalog # AP78797

Product Information

| | |
|--------------------------|---|
| Application | WB, IHC-P, IF, ICC, IP |
| Primary Accession | Q92785 |
| Reactivity | Rat, Human, Mouse |
| Host | Rabbit |
| Clonality | Monoclonal Antibody |
| Isotype | IgG |
| Conjugate | Unconjugated |
| Immunogen | A synthesized peptide derived from human DPF2 |
| Purification | Affinity Chromatography |
| Calculated MW | 44155 |


Additional Information

| | |
|--------------------|--|
| Gene ID | 5977 |
| Other Names | DPF2 |
| Dilution | WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A IP~~N/A |
| Format | Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |

Protein Information

| | |
|--------------------------|---|
| Name | DPF2 |
| Synonyms | BAF45D, REQ, UBID4 |
| Function | Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: 27775714 , PubMed: 28533407). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: 28533407). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: 20460684). |
| Cellular Location | Nucleus. Cytoplasm |
| Tissue Location | Ubiquitous. |

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.