10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

LAG3 Rabbit mAb

Catalog # AP78869

Product Information

Application	IHC-P, IF, ICC
Primary Accession	P18627
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human CD223
Purification	Affinity Chromatography
Calculated MW	57449

Additional Information

Gene ID	3902
Other Names	LAG3
Dilution	IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	LAG3 {ECO:0000303|PubMed:35761082, ECO:0000312|HGNC:HGNC:6476}
Function	[Lymphocyte activation gene 3 protein]: Inhibitory receptor on antigen activated T-cells (PubMed:20421648, PubMed:35761082, PubMed:7805750, PubMed:8647185). Delivers inhibitory signals upon binding to ligands, such as MHC class II, its main ligand present at the surface of antigen-presenting cells (APCs), and FGL1, which is secreted by hepatocytes and certain types of tumor cells (PubMed:30580966, PubMed:32920841, PubMed:35761082, PubMed:39671469, PubMed:7589152, PubMed:8647185, PubMed:9159144). Ligand-binding initiates a signaling that inhibits the T-cell receptor (TCR) in the immunological synapse, preventing T-cell activation (PubMed:40101708). Mechanistically, ligand-binding promotes (1) ubiquitination of the KIEELE motif, unleashing the RRFSALE motif from the membrane and (2) leading to the formation of condensates with the TCR component CD3E, thereby disrupting the association between CD3E and LCK and preventing TCR activation (PubMed:40101708, PubMed:40592325). May inhibit antigen-

	specific T-cell activation in synergy with PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity).
Cellular Location	[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein. Note=Clusters on the T-cell surface following ligand-binding
Tissue Location	Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.