

## LAG3 Rabbit mAb

Catalog # AP78869

## **Product Information**

Application IHC-P, IF, ICC
Primary Accession P18627
Reactivity Human
Host Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human CD223

**Purification** Affinity Chromatography

Calculated MW 57449

## **Additional Information**

**Gene ID** 3902

Other Names LAG3

**Dilution** IHC-P~~N/A IF~~1:50~200 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

## **Protein Information**

Name LAG3 ( HGNC:6476)

**Synonyms** FDC

**Function** Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen

activated T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:20421648, PubMed:7805750, PubMed:8647185). Also mediates

immune tolerance: constitutively expressed on a subset of regulatory T-cells

(Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:8647185).

**Cellular Location** [Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I

membrane protein

**Tissue Location** Primarily expressed in activated T-cells and a subset of natural killer (NK)

cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.