

TBK1 Antibody (S172)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7887a

Product Information

Application	WB, IHC-P, E
Primary Accession	<u>Q9UHD2</u>
Other Accession	<u>Q6DFJ6</u> , <u>Q9WUN2</u> , <u>NP_037386</u>
Reactivity	Human, Mouse
Predicted	Mouse, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	83642
Antigen Region	150-181

Additional Information

Gene ID	29110
Other Names	Serine/threonine-protein kinase TBK1, NF-kappa-B-activating kinase, T2K, TANK-binding kinase 1, TBK1, NAK
Target/Specificity	This TBK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 150-181 amino acids from human TBK.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TBK1 Antibody (S172) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TBK1 {ECO:0000303 PubMed:10581243, ECO:0000312 HGNC:HGNC:11584}
Function	Serine/threonine kinase that plays an essential role in regulating inflammatory responses to foreign agents (PubMed: <u>10581243</u> , PubMed: <u>11839743</u> , PubMed: <u>12692549</u> , PubMed: <u>12702806</u> , PubMed: <u>14703513</u> , PubMed: <u>15367631</u> , PubMed: <u>15485837</u> ,

PubMed: <u>18583960</u> , PubMed: <u>21138416</u> , PubMed: <u>23453971</u> , PubMed: <u>23453972</u> , PubMed: <u>23746807</u> , PubMed: <u>25636800</u> , PubMed: <u>26611359</u> , PubMed: <u>32404352</u> , PubMed: <u>34363755</u> , PubMed: <u>32298923</u>). Following activation of toll-like receptors by viral or bacterial components, associates with TRAF3 and TANK and phosphorylates interferon regulatory factors (IRFs) IRF3 and IRF7 as well as DDX3X (PubMed: <u>12692549</u> , PubMed: <u>12702806</u> , PubMed: <u>14703513</u> ,
PubMed: <u>15367631</u> , PubMed: <u>18583960</u> , PubMed: <u>25636800</u>). This activity allows subsequent homodimerization and nuclear translocation of the IRFs leading to transcriptional activation of pro-inflammatory and antiviral genes
including IFNA and IFNB (PubMed: <u>12702806</u> , PubMed: <u>15367631</u> , PubMed: <u>25636800</u> , PubMed: <u>32972995</u>). In order to establish such an antiviral
state, TBK1 form several different complexes whose composition depends on the type of cell and cellular stimuli (PubMed: <u>23453971</u> , PubMed: <u>23453972</u> , PubMed: <u>23746807</u>) Plays a key role in IRF3 activation: acts by first
phosphorylating innate adapter proteins MAVS, STING1 and TICAM1 on their pLxIS motif, leading to recruitment of IRF3, thereby licensing IRF3 for phosphorylation by TBK1 (PubMed:25636800, PubMed:30842653
PubMed: <u>37926288</u>). Phosphorylated IRF3 dissociates from the adapter proteins, dimerizes, and then enters the nucleus to induce expression of
interferons (PubMed: <u>25636800</u>). Thus, several scaffolding molecules including FADD, TRADD, MAVS, AZI2, TANK or TBKBP1/SINTBAD can be recruited to the TBK1-containing- complexes (PubMed: <u>21931631</u>). Under particular conditions, functions as a NF-kappa-B effector by phosphorylating NF-kappa-B inhibitor
alpha/NFKBIA, IKBKB or RELA to translocate NF-Kappa-B to the nucleus (PubMed: <u>10783893</u> , PubMed: <u>15489227</u>). Restricts bacterial proliferation by
enhancing LC3 binding affinity and antibacterial autophagy (PubMed: <u>21617041</u>). Phosphorylates SMCR8 component of the
C9orf72-SMCR8 complex, promoting autophagosome maturation (PubMed: <u>27103069</u>). Phosphorylates ATG8 proteins MAP1LC3C and GABARAPL2, thereby preventing their delipidation and premature removal
energy balance regulation by sustaining a state of chronic, low-grade inflammation in obesity, which leads to a negative impact on insulin
the endosomal sorting complex required for transport-I (ESCRT-I) subunit VPS37C (PubMed: <u>21270402</u>). Phosphorylates Borna disease virus (BDV) P protein (PubMed: <u>16155125</u>). Plays an essential role in the TLR3- and IFN-
dependent control of herpes virus HSV-1 and HSV-2 infections in the central nervous system (PubMed:22851595). Acts both as a positive and negative regulator of the mTORC1 complex, depending on the context: activates mTORC1 in response to growth factors by catalyzing phosphorylation of MTOR, while it limits the mTORC1 complex by promoting phosphorylation of
RPTOR (PubMed: <u>29150432</u> , PubMed: <u>31530866</u>). Acts as a positive regulator of the mTORC2 complex by mediating phosphorylation of MTOR, leading to increased phosphorylation and activation of AKT1 (By similarity).
regulation of TNF-induced RIPK1- mediated cell death, probably acting via CYLD phosphorylation that in turn controls RIPK1 ubiquitination status (PubMed: <u>34363755</u>). Also participates in the differentiation of T follicular regulatory cells together with the receptor ICOS (PubMed: <u>27135603</u>).
Cytoplasm. Note=Upon mitogen stimulation or triggering of the immune system, TBK1 is recruited to the exocyst by EXOC2.
and a second

Ubiquitous with higher expression in testis. Expressed in the ganglion cells, nerve fiber layer and microvasculature of the retina.

Cellular Location

Tissue Location

Background

The NF-kappa-B (NFKB) complex of proteins is inhibited by I-kappa-B (IKB) proteins, which inactivate NFKB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IKB proteins by IKB kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation and nuclear translocation of the NFKB complex. TKB is similar to IKB kinases and can mediate NFKB activation in response to certain growth factors. The protein can form a complex with the IKB protein TANK and TRAF2 and release the NFKB inhibition caused by TANK.

References

Deng,W., J. Biol. Chem. 283 (51), 35590-35597 (2008) Chessler,A.D., J. Immunol. 181 (11), 7917-7924 (2008) Soulat,D., EMBO J. 27 (15), 2135-2146 (2008)

Images



AP7887a staining TBK1 in human brain tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-TBK1 Antibody (S172) at 1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: HT-1080 whole cell lysate Lane 3: K562 whole cell lysate Lane 4: mouse testis lysate Lane 5: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size :83kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

• <u>Capsular Polysaccharide is a Main Component of Mycoplasma ovipneumoniae in the Pathogen-Induced Toll-Like</u> <u>Receptor-Mediated Inflammatory Responses in Sheep Airway Epithelial Cells.</u>

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.