

Phospho-LRP6 (Ser1490) Rabbit mAb

Catalog # AP78927

Product Information

Application	WB
Primary Accession	O75581
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-LRP6 (S1490)
Purification	Affinity Chromatography
Calculated MW	180429

Additional Information

Gene ID	4040
Other Names	LRP6
Dilution	WB~1/500-1/1000
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	LRP6
Function	Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes (PubMed: 11357136 , PubMed: 11448771 , PubMed: 15778503 , PubMed: 16341017 , PubMed: 16513652 , PubMed: 17326769 , PubMed: 17400545 , PubMed: 19107203 , PubMed: 19293931 , PubMed: 19801552 , PubMed: 28341812 , PubMed: 34896607). Cell-surface coreceptor of Wnt/beta-catenin signaling, which plays a pivotal role in various processes including retinal angiogenesis and bone formation (PubMed: 11357136 , PubMed: 11448771 , PubMed: 15778503 , PubMed: 16341017 , PubMed: 16513652 , PubMed: 17326769 , PubMed: 17400545 , PubMed: 19107203 , PubMed: 19293931 , PubMed: 19801552 , PubMed: 28341812 , PubMed: 34896607). The Wnt-induced Fzd/LRP6 coreceptor complex recruits DVL1 polymers to the plasma membrane which, in turn, recruits the

AXIN1/GSK3B-complex to the cell surface promoting the formation of signalosomes and inhibiting AXIN1/GSK3-mediated phosphorylation and destruction of beta-catenin (PubMed:[16513652](#)). Required for posterior patterning of the epiblast during gastrulation (By similarity).


Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum. Membrane raft. Note=On Wnt signaling, undergoes a cycle of caveolin- or clathrin-mediated endocytosis and plasma membrane location. Released from the endoplasmic reticulum on palmitoylation Mono-ubiquitination retains it in the endoplasmic reticulum in the absence of palmitoylation. On Wnt signaling, phosphorylated, aggregates and colocalizes with AXIN1 and GSK3B at the plasma membrane in LRP6- signalosomes (By similarity). Chaperoned to the plasma membrane by HSP90B1 and MESD (PubMed:23572575). {ECO:0000250 | UniProtKB:O88572, ECO:0000269 | PubMed:23572575}

Tissue Location

Widely coexpressed with LRP5 during embryogenesis and in adult tissues

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.