

# PI3 Kinase p110 alpha Rabbit mAb

Catalog # AP78949

#### **Product Information**

ApplicationWB, IP, ICCPrimary AccessionP42336ReactivityHumanHostRabbit

**Clonality** Monoclonal Antibody

Calculated MW 124284

#### **Additional Information**

**Gene ID** 5290

Other Names PIK3CA

**Dilution** WB~~1/500-1/1000 IP~~N/A ICC~~N/A

Format 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide

and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

### **Protein Information**

Name PIK3CA

**Function** Phosphoinositide-3-kinase (PI3K) phosphorylates phosphatidylinositol (PI)

and its phosphorylated derivatives at position 3 of the inositol ring to produce


3-phosphoinositides (PubMed: 15135396, PubMed: 23936502,

PubMed:<u>28676499</u>). Uses ATP and PtdIns(4,5)P2 (phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3) (PubMed:<u>15135396</u>, PubMed:<u>28676499</u>). PIP3 plays a key role by recruiting PH domain- containing proteins to the membrane, including AKT1 and PDPK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors. Involved in the activation of AKT1 upon stimulation by receptor tyrosine kinases ligands such as EGF, insulin, IGF1, VEGFA and PDGF. Involved in signaling via insulin-receptor substrate (IRS) proteins. Essential in endothelial cell migration during vascular development through VEGFA signaling, possibly by regulating RhoA activity. Required for lymphatic

vasculature development, possibly by binding to RAS and by activation by EGF and FGF2, but not by PDGF. Regulates invadopodia formation through the PDPK1-AKT1 pathway. Participates in cardiomyogenesis in embryonic stem cells through a AKT1 pathway. Participates in vasculogenesis in embryonic

stem cells through PDK1 and protein kinase C pathway. In addition to its lipid kinase activity, it displays a serine-protein kinase activity that results in the autophosphorylation of the p85alpha regulatory subunit as well as phosphorylation of other proteins such as 4EBP1, H-Ras, the IL-3 beta c receptor and possibly others (PubMed:23936502, PubMed:28676499). Plays a role in the positive regulation of phagocytosis and pinocytosis (By similarity).

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.