

# ULK1 Rabbit mAb

Catalog # AP79004

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC
<b>Primary Accession</b>	<a href="#">O75385</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Immunogen</b>	A synthesized peptide derived from human ULK1
<b>Purification</b>	Affinity Chromatography
<b>Calculated MW</b>	112631

## Additional Information

---

<b>Gene ID</b>	8408
<b>Other Names</b>	ULK1
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
<b>Format</b>	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	ULK1 {ECO:0000303   PubMed:9693035, ECO:0000312   HGNC:HGNC:12558}
<b>Function</b>	Serine/threonine-protein kinase involved in autophagy in response to starvation (PubMed: <a href="#">18936157</a> , PubMed: <a href="#">21460634</a> , PubMed: <a href="#">21795849</a> , PubMed: <a href="#">23524951</a> , PubMed: <a href="#">25040165</a> , PubMed: <a href="#">29487085</a> , PubMed: <a href="#">31123703</a> ). Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagophores, the precursors of autophagosomes (PubMed: <a href="#">18936157</a> , PubMed: <a href="#">21460634</a> , PubMed: <a href="#">21795849</a> , PubMed: <a href="#">25040165</a> , PubMed: <a href="#">39384743</a> ). Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1) via interaction with RPTOR (PubMed: <a href="#">21795849</a> ). Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity (PubMed: <a href="#">21460634</a> ). May phosphorylate ATG13/KIAA0652 and RPTOR; however such data need additional evidences (PubMed: <a href="#">18936157</a> ). Plays a

role early in neuronal differentiation and is required for granule cell axon formation (PubMed:[11146101](#)). Also phosphorylates SESN2 and SQSTM1 to regulate autophagy (PubMed:[25040165](#), PubMed:[37306101](#)). Phosphorylates FLCN, promoting autophagy (PubMed:[25126726](#)). Phosphorylates AMBRA1 in response to autophagy induction, releasing AMBRA1 from the cytoskeletal docking site to induce autophagosome nucleation (PubMed:[20921139](#)). Phosphorylates ATG4B, leading to inhibit autophagy by decreasing both proteolytic activation and delipidation activities of ATG4B (PubMed:[28821708](#)).

#### Cellular Location


Cytoplasm, cytosol. Preautophagosomal structure. Note=Under starvation conditions, is localized to punctate structures primarily representing the isolation membrane that sequesters a portion of the cytoplasm resulting in the formation of an autophagosome.

#### Tissue Location

Ubiquitously expressed. Detected in the following adult tissues: skeletal muscle, heart, pancreas, brain, placenta, liver, kidney, and lung

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.