

# MEK2 (MAP2K2) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7961a

## Product Information

---

<b>Application</b>	IF, WB, E
<b>Primary Accession</b>	<a href="#">P36507</a>
<b>Other Accession</b>	<a href="#">P36506</a> , <a href="#">Q63932</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	44424
<b>Antigen Region</b>	1-30

## Additional Information

---

<b>Gene ID</b>	5605
<b>Other Names</b>	Dual specificity mitogen-activated protein kinase kinase 2, MAP kinase kinase 2, MAPKK 2, ERK activator kinase 2, MAPK/ERK kinase 2, MEK 2, MAP2K2, MEK2, MKK2, PRKMK2
<b>Target/Specificity</b>	This MEK2 (MAP2K2) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human MEK2 (MAP2K2).
<b>Dilution</b>	IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	MEK2 (MAP2K2) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	MAP2K2
<b>Synonyms</b>	MEK2, MKK2, PRKMK2
<b>Function</b>	Catalyzes the concomitant phosphorylation of a threonine and a tyrosine

residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases (By similarity). Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (PubMed:[29433126](#)).

#### Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Note=Membrane localization is probably regulated by its interaction with KSR1.


## Background

MAP2K2 is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. The inhibition or degradation of this kinase is found to be involved in the pathogenesis of Yersinia and anthrax.


## References

- Burroughs, K.D., et al., Mol. Cancer Res. 1(4):312-322 (2003).  
Tran, H., et al., Mol. Cell. Biol. 23(20):7177-7188 (2003).  
Li, S.P., et al., Cancer Res. 63(13):3473-3477 (2003).  
Li, Y., et al., J. Biol. Chem. 278(16):13663-13671 (2003).  
Liu, X., et al., J. Biol. Chem. 277(42):39312-39319 (2002).

## Images



Fluorescent confocal image of U251 cell stained with MEK2 (MAP2K2) Antibody (N-term)(Cat#AP7961a). U251 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with MEK2 (MAP2K2) primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). MEK2 (MAP2K2) immunoreactivity is localized to Cytoplasm significantly.



Western blot analysis of anti-MAP2K2 Pab (Cat. #AP7961a) in HL-60 cell lysate. MAP2K2 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.