10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ULK3 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8115a

Product Information

Application	WB, IHC-P, E
Primary Accession	Q6PHR2
Other Accession	Q3U3Q1
Reactivity	Human, Rat, Mouse
Predicted	Mouse, Bovine, Canine, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	53444
Antigen Region	14-43

Additional Information

Gene ID	25989
Other Names	Serine/threonine-protein kinase ULK3, Unc-51-like kinase 3, ULK3
Target/Specificity	This ULK3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 14-43 amino acids from the N-terminal region of human ULK3.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ULK3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ULK3 {ECO:0000303|PubMed:19878745, ECO:0000312|HGNC:HGNC:19703}
Function	Serine/threonine protein kinase that acts as a regulator of smoothened signaling and autophagy (PubMed:19279323, PubMed:19878745, PubMed:20643644, PubMed:31279575). Acts as a positive regulator of smoothened signaling in the presence of hedgehog (DHH, IHH or SHH) ligand:

	autophosphorylates and mediates phosphorylation of full-length GLI2, promoting subsequent phosphorylation by CDK1 and GLI2 dissociation from SUFU and translocation to the nucleus (PubMed:19878745, PubMed:20643644, PubMed:31279575). Also phosphorylates as GLI1 and GLI3, although less efficiently (PubMed:19878745, PubMed:20643644). In the absence of hedgehog ligand, interacts with SUFU, thereby inactivating the protein kinase activity and preventing phosphorylation of GLI proteins (GLI1, GLI2 and/or GLI3) (PubMed:20643644). Also acts as a regulator of autophagy: following cellular senescence, able to induce autophagy (PubMed:19279323).
Cellular Location	Cytoplasm. Note=Localizes to pre-autophagosomal structure during cellular senescence.
Tissue Location	Widely expressed. Highest levels observed in fetal brain. In adult tissues, high levels in brain, liver and kidney, moderate levels in testis and adrenal gland and low levels in heart, lung, stomach, thymus, prostate and placenta. In the brain, highest expression in the hippocampus, high levels also detected in the cerebellum, olfactory bulb and optic nerve. In the central nervous system, lowest levels in the spinal cord


Background

ULK3 is a member of the ULK kinase family, one of three human homologs of the yeast autophagy-specific kinase.


References

The MGC Project Team, Genome Res. 14:2121-2127(2004).

Images



Anti-ULK3 Antibody (N-term) at 1:2000 dilution + LNCaP whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human testis tissue reacted with *ULK3 antibody (N-term) (Cat.#AP8115a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.