10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

YWHAB Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8155b

Product Information

Application	IHC-P, WB, IF, E
Primary Accession	P31946
Other Accession	Q5PRD0, P35213, Q9CQV8, Q4R572, Q5ZLQ6, P68250
Reactivity	Human
Predicted	Bovine, Chicken, Monkey, Mouse, Rat, Zebrafish, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB4064
Calculated MW	28082
Antigen Region	217-246

Additional Information

Gene ID	7529
Other Names	14-3-3 protein beta/alpha, Protein 1054, Protein kinase C inhibitor protein 1, KCIP-1, 14-3-3 protein beta/alpha, N-terminally processed, YWHAB
Target/Specificity	This YWHAB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 217-246 amino acids from the C-terminal region of human YWHAB.
Dilution	IHC-P~~1:100~500 WB~~1:1000 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	YWHAB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	YWHAB
Function	Adapter protein implicated in the regulation of a large spectrum of both

general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner. Negative regulator of osteogenesis. Blocks the nuclear translocation of the phosphorylated form (by AKT1) of SRPK2 and antagonizes its stimulatory effect on cyclin D1 expression resulting in blockage of neuronal apoptosis elicited by SRPK2. Negative regulator of signaling cascades that mediate activation of MAP kinases via AKAP13.

Cellular Location Cytoplasm. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV


Background

YWHAB belongs to the 14-3-3 family of proteins, members of which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals. The encoded protein has been shown to interact with RAF1 and CDC25 phosphatases, suggesting that it may play a role in linking mitogenic signaling and the cell cycle machinery. Two transcript variants differing in the 5' UTR, but encoding the same protein, have been identified for the gene. Both variants encode the same protein, however, they are differentially expressed in hematopoietic cells.


References

Komori, T., et al., Acta Neuropathol. 106(1):66-70 (2003). Cavet, M.E., et al., J. Biol. Chem. 278(20):18376-18383 (2003). Li, Y., et al., J. Biol. Chem. 278(16):13663-13671 (2003). Shumway, S.D., et al., J. Biol. Chem. 278(4):2089-2092 (2003). Parvaresch, S., et al., FEBS Lett. 532(3):357-362 (2002).

Images




Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.




The anti-YWHAB Pab (Cat. #AP8155b) is used in Western blot to detect YWHAB in Jurkat cell lysate.

Western blot analysis of YWHAB (arrow) using rabbit polyclonal YWHAB Antibody (C-term) (Cat. #AP8155b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1)



or transiently transfected (Lane 2) with the YWHAB gene.



Confocal immunofluorescent analysis of YWHAB Antibody (C-term)(Cat#AP8155b) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.