

PPM1D Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8437B

Product Information

Application	WB, E
Primary Accession	O15297
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	66675
Antigen Region	571-602

Additional Information

Gene ID	8493
Other Names	Protein phosphatase 1D, Protein phosphatase 2C isoform delta, PP2C-delta, Protein phosphatase magnesium-dependent 1 delta, p53-induced protein phosphatase 1, PPM1D, WIP1
Target/Specificity	This PPM1D antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 571-602 amino acids from the C-terminal region of human PPM1D.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PPM1D Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPM1D
Synonyms	WIP1
Function	Involved in the negative regulation of p53 expression (PubMed: 23242139). Required for the relief of p53-dependent checkpoint mediated cell cycle

arrest. Binds to and dephosphorylates 'Ser-15' of TP53 and 'Ser-345' of CHEK1 which contributes to the functional inactivation of these proteins (PubMed:[15870257](#), PubMed:[16311512](#)). Mediates MAPK14 dephosphorylation and inactivation (PubMed:[21283629](#)). Is also an important regulator of global heterochromatin silencing and critical in maintaining genome integrity (By similarity).

Cellular Location

Nucleus. Cytoplasm, cytosol


Tissue Location

Expressed in fetal and adult brain. Also detected in fetal liver and skeletal muscle, but not in their adult counterparts.

Background

PPM1D is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. Expression of this PPM1D gene is induced in a p53-dependent manner in response to various environmental stresses. While being induced by tumor suppressor protein TP53/p53, this phosphatase negatively regulates the activity of p38 MAP kinase, MAPK/p38, through which it reduces the phosphorylation of p53, and in turn suppresses p53-mediated transcription and apoptosis. This phosphatase thus mediates a feedback regulation of p38-p53 signaling that contributes to growth inhibition and the suppression of stress induced apoptosis. The PPM1D gene is located in a chromosomal region known to be amplified in breast cancer. The amplification of this gene has been detected in both breast cancer cell line and primary breast tumors, which suggests a role of this gene in cancer development.

Images



The anti-PPM1D Pab (Cat. #AP8437b) is used in Western blot to detect PPM1D in 293 cell lysate.

Citations

- [p53-Independent expression of wild-type p53-induced phosphatase 1 \(Wip1\) in methylmethane sulfonate-treated cancer cell lines and human tumors.](#)
- [BRCA1-IRIS overexpression abrogates UV-induced p38MAPK/p53 and promotes proliferation of damaged cells.](#)
- [Oncogenic Wip1 phosphatase is inhibited by miR-16 in the DNA damage signaling pathway.](#)
- [The oncogenic phosphatase WIP1 negatively regulates nucleotide excision repair.](#)
- [Wild-type p53-induced phosphatase 1 dephosphorylates histone variant gamma-H2AX and suppresses DNA double strand break repair.](#)
- [Expression of a homeostatic regulator, Wip1 \(wild-type p53-induced phosphatase\), is temporally induced by c-Jun and p53 in response to UV irradiation.](#)
- [Phosphorylation and degradation of MdmX is inhibited by Wip1 phosphatase in the DNA damage response.](#)
- [The estrogen receptor alpha pathway induces oncogenic Wip1 phosphatase gene expression.](#)